

Outcome measure	Dance			Conventional		
	Median change (IQR)	Range p value	Median change (IQR)	Range p value		
ISWT (m)	47.5 (0.0 to 107.5)	-10 to 130 0.14	-	-	-	-
6MWT (m)	-	-	-	35.0* (22.5 to 55.0)	20 to 60	0.07
CAT	-2.0 (-6.8 to 2.0)	-7 to 2 0.46	1.0 (-7.0 to 5.0)	1.0 (-1.0 to 1.0)	-11 to 6 0.89	0.71
mMRC	0.3 (0 to 0.8)	0 to 1 0.31	0.0 (-1.0 to 1.0)	0.0 (-1.0 to 1.0)	-2 to 1 0.71	0.71
Handgrip (kg)	1.0* (-4.8 to 3.8)	-0.2 to 1.2 0.29	13.8** (-4.0 to 0.5)	-	-	-
HADS anxiety	1.0 (-4.8 to 3.8)	-6 to 4 0.59	-1.0 (-4.0 to 0.5)	-1.0 (-4.0 to 0.5)	-4 to 2 0.22	0.22
HADS Depression	-1.0 (-1.0 to 2.8)	-1 to 4 0.71	-1.0 (-2.0 to 0.5)	-1.0 (-2.0 to 0.5)	-2 to 2 0.33	0.33

Change calculated as post - pre PR values; data are presented as median (IQR) to one decimal place; *p* values are presented to two decimal places; significance accepted at *p* <0.05;
 *one missing data value, **one data value only; CAT = COPD assessment test, HADS = hospital anxiety and depression score, ISWT = incremental shuttle walk test, mMRC = modified medical research council dyspnoea scale, 6MWT = 6-minute walk test.